

Pakistan Veterinary Journal

ISSN: 0253-8318 (PRINT), 2074-7764 (ONLINE) DOI: 10.29261/pakvetj/2024.163

RESEARCH ARTICLE

Applicability of Butterfly Pea Flower Extract as an Alternative Natural Dye in Histopathological Canine Mast Cell Tumor Diagnosis

Meennaree Polkaew¹, Pongsiwa Sotthibandhu², Hassadin Boonsriroj³, Suvarin Pavasutthipaisit³, Vissanu Meeyoo⁴ and Araya Suebkhampet^{2*}

¹Master of Science Program in Animal Biotechnology, Faculty of Veterinary Medicine, Mahanakorn University of Technology, Bangkok 10530, Thailand

²Pre-Clinical Veterinary Science Department, Faculty of Veterinary Medicine, Mahanakorn University of Technology, Bangkok 10530, Thailand

³Department of Veterinary Pathology, Faculty of Veterinary Medicine, Mahanakorn University of Technology, Bangkok 10530, Thailand

⁴Centre for Advanced Materials and Environmental Research, Mahanakorn University of Technology, Bangkok 10530, Thailand

*Corresponding author: araya@mut.ac.th

ARTICLE HISTORY (24-120)

Received:	February 29, 2024	
Revised:	March 27, 2024	
Accepted:	April 4, 2024	
Published online:	April 22, 2024	
Key words:		
Butterfly pea		
Clitoria ternatea		
Dye		
Mast cell tumor		
Histopatholog	У	

ABSTRACT

Mast cell tumors (MCTs) are the most common skin tumors in dogs. Clinical data and cytological examination typically suffice for the initial diagnosis. However, prognosis and appropriate treatments necessitate histopathological examination. Amid global concerns regarding hazardous chemicals, the utilization of natural dyes has garnered increased attention. In this context, we focused on using butterfly pea flower (BPF) extract, rich in anthocyanins, as a dye for canine histopathological MCT staining. The dried petal powder was dissolved in distilled water (DW), filtered to obtain a crude extract, and then subjected to a freeze-drying process to preserve its quality. The BPF-dye was prepared by adding DW and 2.5% NaCl into the dried extract. The paraffin serial sections of 32 diagnosed MCTs were stained with the dye for 2 hours. Additionally, the sections were stained with Hematoxylin and Eosin, and Toluidine blue (TB) routine dyes for MCT diagnosis. Sections of non-MCT round cell tumors were also stained with BPF-dye. The results revealed pink-reddish specific staining in the neoplastic mast cell granules, with different staining levels corresponding to the degree of cytoplasmic granulation, similar to TB staining. However, TB staining exhibited blue or metachromasia, indicating differences in staining color. Additionally, the BPF-dye could differentiate MCT from non-MCT round-cell tumors, providing an alternative method of differentiation. The dye maintained its staining ability for 60 days when stored at 4°C. Accordingly, it holds potential as an alternative dye for MCT diagnosis, being locally available and eco-friendly. Further study should focus on improving the stability and longevity of the dye by preserving its anthocyanin content with natural co-pigments for practical

To Cite This Article: Polkaew M, Sotthibandhu P, Boonsriroj H, Pavasutthipaisit S, Meeyoo V and Suebkhampet A, 2024. Applicability of butterfly pea flower extract as an alternative natural dye in histopathological canine mast cell tumor diagnosis. Pak Vet J, 44(2): 266-273. <u>http://dx.doi.org/10.29261/pakvetj/2024.163</u>

INTRODUCTION

application.

Mast cell tumors (MCTs) are the most common skin tumors in dogs (De Nardi *et al.*, 2022). Although they typically form skin nodules, they also affect other organs such as the liver, spleen, and gastrointestinal tract with a low incidence (Pakhrin *et al.*, 2007). Older individuals of certain breeds, including Boxer, Bulldog, and Labrador are susceptible to MCT development (Shoop *et al.*, 2015; Śmiech *et al.*, 2019). A wide variability of MCT biological behaviors, such as variations in anatomical location, the presence of lymph nodes and distant metastases and the occurrence of paraneoplastic syndromes, complicate clinical management for veterinarians and challenge their ability to provide accurate prognoses to pet owners. This can be assessed by clinical staging and histopathological grading (Kiupel and Camus, 2019). Histopathological grading is commonly used in canine cutaneous MCTs including Patnaik and Kiupel systems. The 3-tier grading system of Patnaik et al. (1984) classifies MCTs based on histopathological features as grade I (monomorphic welldifferentiated mast cells), grade II (moderately pleomorphic mast cells), and grade III (poorly differentiated mast cells). However, the novel Kiupel (2tier) grading system includes high-grade MCT that exhibit any of the following criteria: at least 7 mitotic figures in 10 high-power fields (hpf); at least 3 multinucleated cells in 10 hpf; at least 3 bizarre nuclei in 10 hpf; and karvomegalv. The absence of those characteristics indicates low-grade MCT. High-grade tumors were significantly associated with a poor prognosis and shorter survival times (Kiupel et al., 2011). The samples used in this study underwent assessment based on the Kiupel system. To enhance the prognosis accuracy, it is recommended to incorporate immunohistochemistry or molecular techniques in the MCT biological behavior evaluation. This could entail examining factors like proliferation activity or the mutation status of c-Kit (Sabattini et al., 2015; Gobbo et al., 2021).

Hematoxylin and eosin (H&E) are frequently utilized for histopathological staining. Nevertheless, toluidine blue (TB) or the Romanowsky-type dye such as Giemsa's stain and Wright's stain are recommended for identifying mast cells, as they exhibit metachromasia in their granules (Ribatti, 2018). Most of the dyes currently employed are synthetic, despite their effectiveness. However, these synthetic dyes are expensive, present environmental hazards, and are harmful to health, particularly for laboratory personnel (Ratna and Padhi, 2012) Consequently, there has been increasing attention toward the study of natural dyes as an alternative. Plant anthocyanins, found in various flowers and fruits, are potential candidates for natural dyes due to their vivid blue, red, and purple colors (Khoo et al., 2017). It belongs to flavonoids, similar to hematin, a natural pigment found in hematoxylin (Dapson et al., 2010). Numerous studies have demonstrated the utilization of anthocyanin-rich plants in histological staining protocols (Jasphin et al., 2021). However, butterfly pea flowers (BPF) have received limited attention.

Butterfly pea (Clirotia ternatea L.) is indigenous to tropical equatorial regions (Hasanah et al., 2023). Its various parts have been used in traditional medicine due to richness in biologically active compounds its (Chakraborthy et al., 2018; Dhangar et al., 2023). It is now widely cultivated due to the growing popularity of butterfly pea tea, a traditional herb utilized in Southeast Asia as a healthful beverage (Sofyan et al., 2021). BPF-extract is employed as natural food colorants (Roy and Rhim, 2021: Noola et al., 2022). The extract is abundant in ternatin anthocyanins, a group of delphinine glycosides known for their water solubility. These anthocyanins possess a natural pH indicator property, wherein they exhibit a red color in acidic solutions, transition to purple in neutral solutions, and turn blue to greenish-yellow in alkaline solutions (Saptarini et al., 2015; Thuy et al., 2021). Although extensive studies have been conducted on the use of BPFextract in food manufacturing (Shirodkar et al., 2023), there is limited research regarding its application for tissue staining. Our group has studied the utilization of BPF-

extract in animal blood smears and cytological canine MCT staining, which can differentiate blood cell types and respectively identify MCTs. (Suebkhampet and Sotthibandhu, 2012; Suebkhampet et al., 2020). Although most MCTs are easily diagnosed with cytological evaluation, histopathological grading remains crucial. This contributes further diagnostic or prognostic details, assisting in precise treatments. The present study focuses on investigating the staining ability of BPF-extract in histopathological canine MCTs. The longevity of the dve extract after storage was also investigated to determine its potential for further practical application.

MATERIALS AND METHODS

Aqueous extraction of butterfly pea flowers (BPF): The fresh flowers were purchased from Talaad Thai, the wholesale market for agricultural products in Thailand. The petals were air-dried overnight, heat-dried (60°C) for 5 hours, and finely ground using a blender. The petal powder was dissolved in distilled water (DW) in a 1:10 ratio (Suebkhampet and Sotthibandhu, 2012). Then, the petal powder solvent was centrifuged for 2 hours at 56°C. The aqueous extract was press-filtered through gauze in filter cloth on a plastic filter funnel and subsequently filtered using Whatman grade 1 qualitative filter paper on a glass funnel.

Freeze-drying process of BPF-extract: The process was implied to preserve the quality and increase the extract concentration. The filtered extract in a glass beaker was covered with punctured aluminum foil (for light protection and sublimation during freeze drying), and frozen at -40°C)Sanden, SNQ-0205P, Thailand(for 24 hours. Then, it was rapidly moved to the freeze dryer at -107°C (Labogene Scanvac CoolSafe, FRE4502, Denmark) for 72 hours and stored at -20°C until used.

Preparation of the BPF-dye: The BPF-dye was prepared by mixing DW and freeze-dried extract (1:5), followed by adding 2.5% NaCl as a mordant. The dye solution was filtered with Whatman grade 1 qualitative filter paper, and the pH was measured using a CyberScan 1000 pH meter (Eutech Instruments Pte Ltd, Singapore). The pH was adjusted to 1.5 using phosphoric acid. This freshly prepared dye solution was classified as day 0 of BPF-dye preparation.

Measurement of total anthocyanin content: Total anthocyanin of the fresh extract and BPF-dye were measured by a pH-differential method using a UV-Vis spectrophotometer (Mapada, V1100D, United States). This was conducted based on the method described by Lee *et al.* (2005). The samples were diluted in KCl (0.025M, pH 1.0) and CH₃CO₂Na (0.4 M, pH 4.5) buffer solutions (1:20), respectively. The absorbance was measured at 520 nm and 700 nm wavelengths. The obtained values were used to calculate total anthocyanin content, expressed as cyanidine-3-glucoside equivalents in mg/L, as follows:

Anthocyanin content (mg/L) = $A \times MW \times DF \times 10^3$

Where A) = $A_{520nm} - A_{700nm}$ (pH 1.0) $-A_{520nm} - A_{700nm}$ (pH 4.5; MW) molecular weight = (449.2 g/mol for cyanidin-3-glucoside; DF = dilution factor; l= pathlength in cm; ε = 26900 molar extinction coefficients, in L × mol⁻¹ × cm⁻¹ for cyanidin-3-glucoside; and 10^3 = factor for conversion from g to mg.

Histological samples: The paraffin-embedded samples of 32 diagnosed canine cutaneous MCTs, aged 6 to 17 (10.3 \pm 2.8; mean \pm SD), were obtained from the Vet and Vitro Central Laboratory and Animal Diagnostic Laboratory Center (ADLC), Mahanakorn University of Technology, Bangkok, Thailand. They were classified according to the Kiupel grading system. Excerpts of clinical data and tumor grades were summarized in Table 1. Non-MCT round cell tumors; histiocytoma, lymphoma (n=2), plasmacytoma, and transmissible venereal tumors (TVT) (n=3), were included to compare with MCTs. All samples were prepared as paraffin serial sections (2 μ m of thickness).

 Table I: Clinical data and histological grade regarding the thirty-two dogs with cutaneous MCTs.

Variable	Dogs (n =32)
Sex	
Female	15
Male	11
Unknown	6
Breed	
Thai Ridgeback	8
French Bulldog, Pit Bull, Pug	4
Labrador and Golden Retriever	4
Bang Kaew	3
Shih-Tzu	2
Poodle, Chihuahua	2
Crossbreed	3
Unknown	6
Age (years)	
Adults (2-8)	8
Elderly (>8)	16
Unknown	8
Location	
Limb	8
Head, neck, shoulder	4
Mammary gland	3
Genital area	3
Inguinal area	2
Axilla, frank	2
Skin (not indicate area)	7
Histological grade	
Low-grade	14
High-grade	18

Staining process: The sections were deparaffinized in xylene, and rehydrated through a graded isopropanol concentration (100%, 95% and 70%), and then rinsed in DW. Serial sections were separately stained with H&E, TB and BPF-dye, respectively. The slides were dripping with 200 μ l of the BPF-dye for 2 hours in a moist chamber at room temperature (RT). The sections were quickly washed in DW twice, air-dried (RT), dipped in xylene, and mounted using a Permount mounting medium. H&E (Leica Biosystems, Hematoxylin 560 MX, and alcoholic Eosin Y 515, Germany) and TB (Sigma-Aldrich, O T360-5G, Germany) staining were conducted according to the manufacturer's instruction. Sections of non-MCT round cell tumors were stained with BPF-dye as a negative control.

The MCT sections were also stained with BPF-dye and stored for 30 and 60 days (at 4°C) to evaluate the durability of the dye. All stained slides were observed and imaged using a light microscope (ZEISS, Primostar3, Germany) and Labscope version 3.3.1 software.

RESULTS

Total anthocyanin content: Total anthocyanin detected in fresh extract and BPF-dye were 236.46 mg/L and 370.72 mg/L, respectively. The increased anthocyanin content in BPF-dye probably resulted from blending freeze-dried extract with DW in a ratio, which was decreased from 1:10 in the fresh extract to 1:5 in BPF-dye. This enhanced the efficiency of the staining.

BPF-dye staining: The staining results revealed that the BFP dye specifically stained the neoplastic mast cell granules at different levels, varying with the degree of cytoplasmic granulation. It was similar to the TB staining pattern; however, it had a distinct color. BPF-dye staining revealed a pink-reddish color in the granules, while TB staining resulted in a blue or metachromatic appearance. Faintly pink staining was also detected in the tumoral nuclei and collagenous stroma. The same stained areas were observed in MCT serial sections stained with H&E, TB, and BPF-dye, respectively, indicating the specificity of the BPF-dye staining (Fig. 1). Both low-grade and highgrade MCTs with well-differentiated tumoral cells revealed numerous tumor cells stained with BPF-dye. In contrast, high-grade MCTs with poorly differentiated cells exhibited markedly fewer stained tumor cells with the BPF-dye (Fig. 2). The pattern observed was consistent with TB staining. Infiltration of eosinophils was noted within the tumoral tissues, with their nuclei appearing pale pink when stained with the BPF-dye; nonetheless, their granules remained indistinct. (Fig. 2G-, J-white arrowhead).

With H&E staining, well-differentiated tumor cells displayed many finely basophilic granules (Fig. 1L and Fig. 2F). Tumor cells in low-grade MCT were observed clustered or rows between collagen bundles, exhibiting minor nuclear changes (Fig. 2C). In contrast, high-grade MCT displays moderate cellular pleomorphism. In contrast, high-grade MCT displays poorly differentiated tumor cells with moderate cellular pleomorphism, high mitotic activity, and inconspicuous cytoplasmic granules (Fig. 2R). Although the results indicate that BPF-dye is effective in identifying mast cells, it is not suitable for tumor grading, unlike H&E staining. Nevertheless, it acts as a special dye in conjunction with H&E staining for MCT diagnosis.

The BPF-dye maintained its staining ability for about 60 days when stored at 4°C. On days 30 and 60 of the dye preparation, the staining pattern was similar to that on day 0. However, the intensity was slightly higher, and some dye precipitates were detected in the tissues (Fig. 3A-F). The BPF-dye staining was not detected in histiocytoma, lymphoma, plasmacytoma, and TVT samples. (Fig. 3I, J, K and L).



Fig. I: Low-grade canine MCTs were stained with BPF-dye (A, D, G, I), Toluidine blue (TB) (B, E, H, K), and H&E (C, F, I, L) at different BPF-dye The magnifications. specifically bound to MCT granules, appearing as a pink-reddish color (black arrow). The nuclei and collagenous stroma stained faintly pink. The granules stained blue, or metachromatically with TB staining (white arrows), while finely basophilic granules were detected with H&E staining (black arrowhead). Each dye exhibited a consistent staining pattern within the same area across the serial sections.

DISCUSSION

Different dyes are applied to selectively stain cellular structures. The present study highlights specific staining of the BPF-dye, resulting in a pink-reddish color in the MCT granules. This is possibly due to the binding between the flavylium cations of ternatin anthocyanins in the acidic dye solution and the polyanionic heparins rich in the MCT granules (Ribatti, 2018). As anthocyanins possess the ability to change color with pH (Chu *et al.*, 2016), the acidic dye solution and the microenvironment within the granules affect the coloring. Moreover, anthocyanin aggregations, complexation by organic molecules, and the presence of metal ions have been identified as factors influencing coloration (Quina *et al.*, 2009).

The BPF-dye staining was also detected in the MCT nuclei and tumor stroma. This staining pattern may be attributed to the properties of anthocyanins, which have been documented to exhibit binding capabilities to biological membranes, proteins, and DNA. However, their color was lighter than it was in the granules. This may be attributed to electrostatic interactions between the hydroxyl groups of anthocyanins and the polar groups of these macromolecules (Jaldappagari *et al.*, 2008; Dudek *et al.*, 2023). Eosinophils can also be identified to some extent by their pale pink, bilobed nuclei. They are often seen in MCT due to chemotactic factors released by mast cells (Galietta *et al.*, 2023).

MCT serial sections stained with H&E, TB, and BPFdye revealed consistent staining in the same areas, whereas no staining was observed in non-MCT round cell tumors. These findings affirm the specificity of BPF-dye staining. The staining was highlighted in the MCT granules and showed a similar pattern to TB staining, indicating its potential in identifying mast cells. However, it yielded a different color tone. The staining of MCT with BPF-dye and TB appears to be correlated with the amount and size of the granules and the degree of cellular differentiation. In poorly differentiated tumors, contain diminished concentrations of heparin, which is the main substance to interacts with the dye.



Fig. 2: Low-grade and high-grade canine MCTs were stained with BPFdye (A, D, G, J, M, P), TB (B, E, H, K, N, Q) and H&E (C, F, I, L, O, R). The BPF-dye staining revealed a pinkreddish color in the MCT granules of both grades (black arrows), exhibiting varying intensity corresponding to the degree of cytoplasmic granulation. This revealed the same trend as TB (white arrows). High-grade MCTs with poorly-differentiated cells revealed a few tumor cells stained with both dyes (M, N). In this case, mitotic figures were occasionally observed with H&E stain (black arrowheads). Eosinophils were frequently seen in MCTs (white arrowheads).

Poor staining may indicate low production of the granular contents (Simoes *et al.*, 1994). Unlike H&E staining, neither of them provided sufficient detail of cell and tissue structures for tumor grading. Therefore, the BPF-dye is suitable to be used as a special stain, like TB, for confirming mast cells in addition to the routine H&E staining.

The results appear to be consistent with our previous study on cytological staining using ethanolic BPF-dye (Suebkhampet *et al.*, 2020). However, the number of stained granules in cytological staining was fairly lower compared to histopathological staining using aqueous BPF-dye. Furthermore, no staining has been observed within the MCT nuclei. These variations may arise due to various factors, such as the type of solvent and mordant, dye concentration, pH level, and tissue preparation process all of which influence the staining capability (Jasphin *et al.*, 2023). Some authors have noted that aqueous BPF-extract yields significantly higher anthocyanin content than ethanolic-extract (Vidana Gamage *et al.*, 2021; Netravati *et*

al., 2022), indicating possible benefits in enhancing dyeing efficiency. Considering the advantages of using an aqueous dye, it would be interesting to investigate whether this formula can also effectively stain MCT cytology. Using water as a solvent is both safe and cost-effective.

We included NaCl as a mordant in the BPF-dye similar to TB. Both dyes have pH levels of 1.5 and 2.5, respectively. In contrast, the previous study on MCT cytology used aluminum chloride as a mordant, resulting in a dye with a very low pH of 0.58. Therefore, it is plausible that the similarity in staining patterns observed between the BPF-dye and TB in this study is due to using the same mordant. NaCl is considered safer than other metallic mordants, which are known to be relatively toxic. (Suebkhampet and Naimon, 2014).

Staining remained effective with the BPF-dye, despite being stored at 4°C for 60 days, demonstrating its longevity and potential for reuse. However, we observed dye precipitation in the tissue, prompting the recommendation to filter it with a paper filter before use. Storing the dye at 271





4°C may help minimize anthocyanin degradation due to light and higher temperatures (Vidana Gamage et al., 2023). In addition, the acidic dye condition is suitable for its stabilization (Sachdev et al., 2021). Furthermore, freeze-drying helps to retain the anthocyanin content of the BPF-extract. Anthocyanins may be degraded due to a combination of many factors such as light, temperature, pH, solvents, and co-pigmentation (Hou et al., 2013). Copigmentation with organic acid can improve anthocyanins' stability by protecting them from the nucleophilic addition of water; otherwise, the flavalium ion becomes pseudobase or colorless (Luo et al., 2022). After 6 months of staining, the tissues were also preliminary assessed to evaluate for the BPF-dye fastness. It found that color intensity had decreased compared to the initial staining, with the color shifting from pink-reddish to a dull purple. This may be due to the alteration or degradation of anthocyanins stained on the tissues. However, it remains adequate for identifying mast cells.

Although the BPF-dye demonstrates effective staining, it requires a longer staining duration compared to TB staining. Further enhancement is achievable

through specific developments. For instance, pretreating the dye with natural co-pigments such as catechin or tannin, which are known to bolster anthocyanin stability and color intensity (Khoo et al., 2017; Charurungsipong al., 2020), could be explored. Additionally, et encapsulating the dye using substances like maltodextrin or Arabic gum has proven effective in preserving anthocyanins, with a half-life of 14 to 19 months, in freeze-dried powder during storage (Hamzah et al., 2013; Fuzetti et al., 2022). These approaches may improve the preservation of the BPF-dye's anthocyanin content and enhance staining efficacy, leading to a reduction in staining time and consequently making the staining process more efficient. However, a significant limitation is the unstable consistency of the anthocyanin content in the BPF-dye extract from batch to batch. Factors such as plant cultivation, harvesting season, and plant age significantly affect the anthocyanin content in the extract (Andrzejewska et al., 2015; Dabravolski and Isayenkov, 2023). Therefore, it is imperative to establish a standardized anthocyanin concentration in the BPF-dye to ensure its practical applicability.

Conclusions: The current investigation presents an innovative method for employing the BPF-dye as an alternative dye in histopathological diagnosis of canine cutaneous MCTs. It offers the advantages of specificity and eco-friendliness. Future research should prioritize enhancing the dye's stability and longevity to facilitate practical application. Additionally, this might broaden its applicability to diverse tissues, including visceral MCTs, consequently enriching diagnostic capabilities within veterinary pathology.

Acknowledgments: The authors are thankful to Vet and Vitro Central Laboratory and ADLC, Mahanakorn University of Technology, Bangkok, Thailand for providing the paraffin-embedded samples. This research was financially supported by Mahanakorn University of Technology, Bangkok, Thailand.

Authors contribution: MP, PS, and SP performed experiments. AS provided supervision, developed the initial concept, and wrote the article. AS and PS designed the experiments and interpreted the data. PS made the figures and legends. WM advised on the BPF extraction and freeze-drying process. HB compiled samples, and clinical data and interpreted H&E staining. All the authors critically revised the manuscript.

REFERENCES

- Andrzejewska J, Sadowska K, Klóska L, et al., 2015. The effect of plant age and harvest time on the content of chosen components and antioxidative potential of black chokeberry fruit. Acta Sci Pol Hortorum Cultus 14(4):105-114.
- Chakraborthy GS, Kumar V, Gupta S, et al., 2018. Phytochemical and pharmacological aspects of *Cliteria ternatea*- A review. J Appl Pharm Sci Res 1:3-9.
- Charurungsipong P, Tangduangdee C, Amornraksa S, et al., 2020. Improvement of anthocyanin stability in Butterfly pea flower extract by co-pigmentation with catechin. E3S Web of Conf 141:1-4.
- Chu B-S, Wilkin J, House M, et al., 2016. Effect of sucrose on thermal and pH stability of Clitoria ternatea extract. Int J Food Process Technol 3(1):11-17.
- Dabravolski SA and Isayenkov SV, 2023. The role of anthocyanins in plant tolerance to drought and salt stresses. Plants 12(13):2558.
- Dapson R, Horobin RW and Kiernan J, 2010. Hematoxylin shortages: their causes and duration, and other dyes that can replace hemalum in routine hematoxylin and eosin staining. Biotechnic Histochem 85(1):55-63.
- De Nardi AB, Dos Santos Horta R, Fonseca-Alves CE, et al., 2022. Review diagnosis, prognosis and treatment of canine cutaneous and subcutaneous mast cell tumors. Cells 11(4):618.
- Dhangar PD, Shimpi H, Newadkar R, et al., 2023. A review on butterfly pea: An emerging plant with applications in food and cosmetics. Int Res J Mod Eng Technol Sci 5(5):1186-1191.
- Dudek A, Strugała-Danak P, Kral T, et al., 2023. An analysis of interactions between three structurally diverse anthocyanidins, as well as their glucosides, and model biological membranes, albumin, and plasmid DNA. Sci Rep 13(1):12883.
- Fuzetti CG, de Castilhos MBM and Nicoletti VR, 2022. Microencapsulation of natural blue dye from butterfly pea (*Clitoria ternatea L.*) flowers: The application of different carriers. J Food Process Preserv 46(4):16420.
- Galietta V, Parisi S, Cocumelli C, et al., 2023. Preliminary assessment of Tumor-Associated Tissue Eosinophilia (TATE) in canine mast cell tumors: Prevalence and prognostic relevance and its association with neoangiogenesis. Animals 13(2):283.
- Gobbo F, Sarli G, De Silva M, et al., 2021. A double histochemical/immunohistochemical staining for the identification of canine mast cells in light microscopy. Vet Sci 8(10):299.
- Hamzah Y, Jumat NA and Sembok W, 2013. Effect of drying on the storage stability of encapsulated anthocyanins powder extract from

Butterfly pea flower (*Clitoria ternatea*). 13th ASEAN Food Conference, Singapore; 9-11 Sept, 2013, pp:1-10.

- Hasanah NN, Azman EM, Rozzamri A, et al., 2023. A systematic review of butterfly pea flower (*Clitoria ternatea* L.): Extraction and application as a food freshness pH-Indicator for polymer-based intelligent packaging. Polym 15:2541:1-20.
- Hou Z, Qin P, Zhang Y, et al., 2013. Identification of anthocyanins isolated from black rice (*Oryza sativa* L.) and their degradation kinetics. Food Res Int 50(2):691-697.
- Jaldappagari S, Motohashi N, Puttagiddappa MG, et al., 2008. Bioactive mechanism of interaction between anthocyanins and macromolecules like DNA and proteins. Top Heterocycl Chem 15:49–65.
- Jasphin S, Raghavendra AP, Solomon MC, et al., 2023. Anthocyanins: A hue for histology-systematic review. MJAS 42(1):106-125.
- Khoo HE, Azlan Ä, Tang ST, *et al.*, 2017. Anthocyanidins and anthocyanins: colored pigments as food, pharmaceutical ingredients, and the potential health benefits. Food Nutr Res 61:1-21.
- Kiupel M and Camus M, 2019. Diagnosis and prognosis of canine cutaneous mast cell tumors. Vet Clin North Am Small Anim Pract 49(5):819–836.
- Kiupel M, Webster JD, Bailey KL, et al., 2011. Proposal of a 2-tier histologic grading system for canine cutaneous mast cell tumors to more accurately predict biological behavior. Vet Pathol 48:147–55.
- Lee J, Durst RW and Wrolstad RE, 2005. Determination of total monomeric anthocyanin pigment content of fruit juices, beverages, natural colorants, and wines by the pH differential method: collaborative study. J AOAC Int 8(5):1269-1278.
- Luo X, Wang R, Wang J, et al., 2022. Acylation of anthocyanins and their applications in the food industry: mechanisms and recent research advances. Foods 11:2166.
- Netravati, Gomez S, Pathrose B, et al., 2022. Comparative evaluation of anthocyanin pigment yield and its attributes from Butterfly pea (*Clitorea ternatea* L.) flowers as prospective food colorant using different extraction methods. Future Foods 6:100199.
- Noola N, Gomez S, Pathrose B, et al., 2022. Comparative evaluation of anthocyanin pigment yield and its attributes from Butterfly pea (*Clitorea ternatea* L.) flowers as prospective food colorant using different extraction methods. Future Foods 6(2):100199.
- Pakhrin B, Kang MS, Bae IH, et al., 2007. Retrospective study of canine cutaneous tumors in Korea J Vet Sci 8(3):229-236.
- Patnaik AK, Ehler WJ and MacEwen EG, 1984. Canine cutaneous mast cell tumor: morphologic grading and survival time in 83 dogs. Vet Pathol 21(5):469-474.
- Quina FH, Moreira PF Jr, Vautier-Giongo C, et al., 2009. Photochemistry of anthocyanins and their biological role in plant tissues. Pure Appl Chem 81:1687-1694.
- Ratna and Padhi BS, 2012. Pollution due to synthetic dyes toxicity & carcinogenicity studies and remediation. Int | Environ Sci 3:940-955.
- Ribatti D, 2018. The staining of mast cells: A historical overview. Int Arch Allergy Immunol 176(1):55-60.
- Roy and Rhim, 2021. Anthocyanin food colorant and its application in pHresponsive color change indicator films. Crit Rev Food Sci Nutr 61(14):2297-2325.
- Sabattini S, Scarpa F, Berlato D, et al., 2015. Histologic Grading of Canine Mast Cell Tumor: Is 2 Better Than 3? Vet Pathol 52(1):70-73.
- Sachdev SS, Chettiankandy TJ, Sonawane SG, et al., 2021. Toward developing natural histologic stains using anthocyanins: A novel approach. J Oral Maxillofac Pathol 25(1):199.
- Saptarini NM, Suryasaputra D and Nurmalia H, 2015. Application of Butterfly Pea (*Clitoria ternatea* Linn) extract as an indicator of acidbase titration. | Chem Pharm Res 7(2):275-280.
- Shirodkar SM, Multisona RR and Gramza-Michalowska A, 2023. The potential for the implementation of pea flower (*Clitoria ternatea*) health properties in food matrix. Appl Sci 13(12):7141.
- Shoop SJW, Marlow S, Church DB, et al., 2015. Prevalence and risk factors for mast cell tumours in dogs in England 2015. Canine Genet Epidemiol 2(1):1-10.
- Simoes JP, Schoning P and Butine M, 1994. Canine mast cell tumors: A comparison of staining techniques. J Vet Diagn Invest 6(4):458–465.
- Śmiech A, Łopuszyński W, Ślaska B, et al., 2019. Occurrence and distribution of canine cutaneous mast cell tumour characteristics among predisposed breeds. 2019. J Vet Res 63:141-148.
- Sofyan J, Permana T and Marpaung AM, 2021. The study of several applicable treatments for serving butterfly pea flower drinks. Adv Biol Sci Res 16:50-56.
- Suebkhampet A and Naimon, 2014. Using dye plant extract for histological staining. J Mahanakorn Vet Med 9(1):63-78.

- Suebkhampet A and Sotthibandhu P, 2012. Effect of using aqueous crude extract from butterfly pea flowers (*Clitoria ternatea* L.) as a dye on animal blood smear staining. Suranaree J Sci Technol 19:15-19.
- Suebkhampet A, Sotthibandhu P and Laoharatchatathanin T, 2020. Butterfly pea flower extract as an alternative dye in cytological canine mast cell tumor staining. Thai J Vet Med 50(4):473-478.
- Thuy NM, Minh VQ, Ben TC, et al., 2021. Identification of anthocyanin compounds in Butterfly pea flowers (*Clitoria ternatea* L.) by

ultraperformance liquid chromatography/ultraviolet coupled to mass spectrometry. Molecules 26(15):4539.

- Vidana Gamage GC, Lim YY and Choo WS, 2021. Anthocyanins from *Clitoria ternatea* flower: Biosynthesis, extraction, stability, antioxidant activity, and applications. Front Plant Sci 12:792303.
- Vidana Gamage GC, Goh JK and Choo WS, 2023. Natural blue colourant preparations from blue pea flower and spirulina: A comparison stability study. Food Chem Adv 3:100457.